

# Extreme IO

Dr Thierry de Baère



Dr Frédéric Deschamps



- 4 year old , 16 kg
- Hepatoblastoma Stage IV + 2 lung metastases

Beyond the evidence  
with 4 year old.....

- 4 year old , 16 kg
- Hepatoblastoma Stage IV + 2 lung metastases
- Received chemotherapy + Surgery for lung metastases



Lung metastases from hepatoblastoma at pathology

**CT before liver resection + 12 new lung metastases**

- 4 year old , 16 kg
- Hepatoblastoma Stage IV + 2 lung metastases
- Received chemotherapy + Surgery for lung metastases
- **+ 12 new lung metastases**
- carboplatin –doxorubicin, changed for Vincristin-Cisplatin-CPT11

>>>> Stable Disease at 6 months

- Multidisciplinary staff:
  - Resection liver tumour
  - After complete resection of lung mets

Secretion of Hepatic Growth Factors after surgery

increases size of the metastases ?




von Schweinitz D. Eur J Pediatr Surg. 1998

von Schweinitz D. Int J Cancer. 2000


- 4 year old , 16 kg
- Hepatoblastoma Stage IV + 2 lung metastases
- Received chemotherapy + Surgery for lung metastases
- **+ 12 new lung metastases**



- 4 mets in right lung
- 8 mets left lung



RFA prefered to surgery







- Minimal invasive technique >>> short delay to liver resection
- 3-8 mm in diameter (high efficacy)
























6 minutes (2' 20W, 2' 40 W, 2' 60W)







Penumbra of ground glass opacification around the tumour

















6 minutes (2' 20W, 2' 40 W, 2' 60W)














6 minutes (2' 20W, 2' 40 W, 2' 60W)




Penumbra of ground glass opacification around the tumour




M 4 20101505  
DoB: Aug 03  
Ex:Jan 19



First session: 8 mets treated in the left lung (2h30)



D1




Delayed pneumothorax D12: asymptomatic/ no treatment


Second session (4 weeks): 4 mets treated in the right lung




Second session (4 weeks): 4 mets treated in the right lung




Second session (4 weeks): 4 mets treated in the right lung





Second session (4 weeks): 4 mets treated in the right lung



Second session (4 weeks): 4 mets treated in the right lung















Day 2: discharged

Day 7: R0 hepatic resection







P 73 2 months


- Complications:

- 1 delayed non-pathological fracture of the rib (at 11 months)
- Asymptomatic
- biopsied



- No other complication

1 new lung mets  
retreated after 8 months  
1 new lung mets  
retreated after 3.5 years



## RFA benefit

- has avoided bilateral thoracotomies
- has allowed surgical liver resection

## RFA benefit

- has avoided bilateral thoracotomies
- has allowed surgical liver resection
- but...
  - no biopsy (too small)
  - imaging follow-up more difficult
  - no data in the literature

Patient is 10 years old and disease free,

we have not seen him since 11 months

Thank you