

For the treatment of attention-deficit/  
hyperactivity disorder (ADHD) in  
children and teens 6 years and older...


What are some  
*reasonable*  
expectations for treatment  
with a nonstimulant?

## When can we expect results?

### Qelbree is a first-of-its-kind, once-a-day nonstimulant treatment approach for ADHD

- **Qelbree** is the first novel nonstimulant approach in ADHD in over a decade

### Qelbree is proven to work in medical studies that were conducted over 6-8 weeks—showing significant reduction in ADHD symptom scores in 4 medical studies of more than 1400 people



- **Qelbree** reduced ADHD symptom scores as early as week 1 for some children

Not everyone responds to medication the same way, and your child's body may need time to adapt. It is important to share how your child is feeling with your healthcare provider (HCP), so treatment can be adjusted if necessary.

#### IMPORTANT SAFETY INFORMATION

Qelbree may increase suicidal thoughts and actions, in children and adults with ADHD, especially within the first few months of treatment or when the dose is changed. Tell your doctor if you or your child have (or if there is a family history of) suicidal thoughts or actions before starting Qelbree. Monitor your or your child's moods, behaviors, thoughts, and feelings during treatment with Qelbree. Report any new or sudden changes in these symptoms right away.

Abbreviation: ADHD, attention-deficit/hyperactivity disorder.

Please see full Important Safety Information on page 4.

## What should we know about safety?

### Qelbree has been proven safe\* in 3 medical studies for the treatment of ADHD in children

- **Qelbree** demonstrated a low incidence of side effects in these studies, with few children or teens stopping treatment with **Qelbree**
- Only 3% of children or teens taking **Qelbree** discontinued taking it in the medical studies due to side effects (compared to 1% taking placebo)

\*Important: Monitor for mood or behavior changes. Your child's HCP may also monitor for changes in blood pressure and heart rate

### Qelbree is a nonstimulant, which means it is not a narcotic or a controlled medication

- There is no evidence of abuse in medical studies

It is important to share how your child is feeling with your HCP, including any side effects. This way, your HCP will better understand what effect your child's treatment may be having, and can adjust treatment if necessary.

#### IMPORTANT SAFETY INFORMATION

The most common side effects of Qelbree in patients 6 to 17 years are sleepiness, not feeling hungry, feeling tired, nausea, vomiting, trouble sleeping, and irritability, and in adults, insomnia, headache, sleepiness, tiredness, nausea, decreased appetite, dry mouth, and constipation. These are not all the possible side effects of Qelbree.



## Can we expect to find it at the pharmacy?

### Qelbree is widely available in pharmacies

Patients that were prescribed Qelbree came from:

**65%** switched to Qelbree from a controlled substance or stimulant medication

**35%** switched to Qelbree from another nonstimulant medication

### Qelbree is backed by a robust savings and support program



- Pay as little as \$20 per prescription each month for a year\* if you have commercial insurance

\*Terms and conditions apply.

If you are having difficulty getting your child's monthly ADHD prescription, ask your HCP if Qelbree is an appropriate treatment option. Your child's HCP may have samples of Qelbree, or can request them from the manufacturer if a trial of Qelbree is an option for your child.

### IMPORTANT SAFETY INFORMATION

**You should not take Qelbree if you or your child:**

take a medicine for depression called a monoamine oxidase inhibitor (MAOI) or have stopped taking an MAOI in the past 14 days. Also, you or your child should avoid alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, and theophylline.


Please see full Important Safety Information on page 4.

## What else can we expect from treatment?



### Qelbree is convenient

- Qelbree is a once-a-day treatment that offers full-day medication coverage
- Qelbree can be prescribed via telehealth; there is no need for a new prescription each month to refill your child's medication
- Qelbree is the only nonstimulant ADHD treatment that can be sprinkled on food to meet the needs of those children or teens unable to swallow capsules
- Qelbree capsule can be swallowed whole (do not cut, crush or chew capsules)
- Qelbree capsule can be opened and sprinkled in its entirety onto a spoonful of pudding or applesauce; the entire mixture needs to be consumed, without chewing, within 15 minutes if mixed with pudding, or within 2 hours if mixed with applesauce; the mixture cannot be stored for future use



Hear what moms have to say about their children's real life experience on Qelbree at [Qelbree.com](https://www.qelbree.com).



### IMPORTANT SAFETY INFORMATION

**Qelbree can increase blood pressure and heart rate.** Your or your child's doctor will monitor these vital signs.

**Qelbree**<sup>®</sup> ONCE-A-DAY  
viloxazine  
extended-release capsules  
100 mg 150 mg 200 mg

## INDICATION

Qelbree is a prescription medicine used to treat ADHD in adults and children 6 years and older.

## IMPORTANT SAFETY INFORMATION

**Qelbree may increase suicidal thoughts and actions, in children and adults with ADHD, especially within the first few months of treatment or when the dose is changed. Tell your doctor if you or your child have (or if there is a family history of) suicidal thoughts or actions before starting Qelbree. Monitor your or your child's moods, behaviors, thoughts, and feelings during treatment with Qelbree. Report any new or sudden changes in these symptoms right away.**

### **You should not take Qelbree if you or your child:**

Take a medicine for depression called a monoamine oxidase inhibitor (MAOI) or have stopped taking an MAOI in the past 14 days. Also, you or your child should avoid alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, and theophylline.

**Qelbree can increase blood pressure and heart rate.** Your or your child's doctor will monitor these vital signs.


**Qelbree may cause manic episodes in patients with bipolar disorder.** Tell your doctor if you or your child show any signs of mania.

**Do not drive or operate heavy machinery until you know how Qelbree will affect you or your child.** Qelbree may cause you or your child to feel sleepy or tired.

**The most common side effects of Qelbree** in patients 6 to 17 years are sleepiness, not feeling hungry, feeling tired, nausea, vomiting, trouble sleeping, and irritability, and in adults, insomnia, headache, sleepiness, tiredness, nausea, decreased appetite, dry mouth, and constipation. These are not all the possible side effects of Qelbree.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see Brief Summary, including Boxed Warning, on pages 6 and 7.



## Real moms share their children's treatment experiences with Qelbree



Scan here to watch them!

**Qelbree**<sup>®</sup> ONCE-A-DAY  
viloxazine  
extended-release capsules  
100 mg 150 mg 200 mg

### INDICATION

Qelbree is a prescription medicine used to treat ADHD in patients 6 years and older.

### IMPORTANT SAFETY INFORMATION

**Qelbree may increase suicidal thoughts and actions, in children and adults with ADHD, especially within the first few months of treatment or when the dose is changed. Tell your doctor if you or your child have (or if there is a family history of) suicidal thoughts or actions before starting Qelbree. Monitor your or your child's moods, behaviors, thoughts, and feelings during treatment with Qelbree. Report any new or sudden changes in these symptoms right away.**

Please see full Important Safety Information on page 4.

<b>Consumer Brief Summary</b>	
<b>Qelbree® (Kel’bree) (viloxazine extended-release capsules), for oral use</b>	
<b>What is the most important information I should know about QELBREE?</b>	<p><b>Qelbree can cause serious side effects, including:</b></p> <ul style="list-style-type: none"> <li>• <b>Increased risk of suicidal thoughts or actions.</b> Qelbree may increase suicidal thoughts or actions in children and adults with attention deficit hyperactivity disorder (ADHD), <b>especially within the first few months of treatment or when the dose is changed.</b></li> </ul> <p><b>How can I watch for and try to prevent suicidal thoughts and actions?</b></p> <ul style="list-style-type: none"> <li>• Pay close attention to any changes, especially new or sudden changes in mood, behavior, thoughts, or feelings or if you or your child develops suicidal thoughts or actions. This is very important when Qelbree treatment is started or when the dose is changed. Contact your healthcare provider right away if you or your child has any sudden changes in mood, behavior, thoughts, or feelings, or develops suicidal thoughts or actions.</li> <li>• Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.</li> </ul> <p><b>Call your healthcare provider or get emergency help right away if you or your child has any of the following symptoms, especially if they are new, worse, or worry you:</b></p> <ul style="list-style-type: none"> <li>• attempts to commit suicide</li> <li>• new or worse depression</li> <li>• feeling very agitated or restless</li> <li>• trouble sleeping (insomnia)</li> <li>• acting aggressive, being angry or violent</li> <li>• an extreme increase in activity and talking (mania)</li> <li>• thoughts about suicide or dying</li> <li>• new or worse anxiety</li> <li>• panic attacks</li> <li>• new or worse irritability</li> <li>• acting on dangerous impulses</li> <li>• other unusual changes in behavior or mood.</li> </ul> <p>See <b>What are the possible side effects of Qelbree?</b> for more information.</p>
<b>What is QELBREE used for?</b>	<p>Qelbree is a prescription medicine used to treat ADHD in adults and children 6 years and older.</p> <p>It is not known if Qelbree is safe and effective in children less than 6 years of age.</p>
<b>Do not take QELBREE if you or your child:</b>	<ul style="list-style-type: none"> <li>• Takes a medicine used to treat depression called a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you or your child takes an MAOI.</li> <li>• Stopped taking an MAOI in the last 14 days.</li> <li>• Takes alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, or theophylline.</li> </ul>
<b>What are the possible side effects of QELBREE?</b>	<p><b>Qelbree can cause serious side effects, including:</b></p> <ul style="list-style-type: none"> <li>• See <b>What is the most important information I should know about Qelbree?</b></li> <li>• <b>Increased blood pressure and heart rate.</b> Your healthcare provider should check you or your child’s blood pressure and heart rate before starting and during treatment with Qelbree.</li> <li>• <b>Manic episodes.</b> Manic episodes may happen in people with bipolar disorder who take Qelbree. Symptoms may include: <ul style="list-style-type: none"> <li>• greatly increased energy</li> <li>• racing thoughts</li> <li>• unusually grand ideas</li> <li>• talking more or faster than usual</li> <li>• severe trouble sleeping</li> <li>• reckless behavior</li> <li>• excessive happiness or irritability</li> </ul> </li> <li>• <b>Sleepiness and tiredness:</b> Qelbree may cause you or your child to feel sleepy or tired. Do not drive or operate heavy machinery until you know how Qelbree will affect you.</li> </ul>
<b>What should I tell my healthcare provider?</b>	<p><b>Before taking Qelbree, tell your healthcare provider about all your or your child’s medical conditions, including if you or your child:</b></p>

	<ul style="list-style-type: none"> <li>• have a family history of suicide, bipolar disorder, depression, mania or hypomania</li> <li>• have blood pressure or heart rate problems</li> <li>• have severe kidney problems. Your healthcare provider may lower the dose of Qelbree.</li> <li>• are pregnant or plan to become pregnant. Qelbree may cause harm to the mother when taken during pregnancy. <ul style="list-style-type: none"> <li>○ Tell your healthcare provider right away if you or your child becomes pregnant or thinks they are pregnant during treatment with Qelbree.</li> <li>○ There is a pregnancy registry for females who are exposed to Qelbree during pregnancy. Talk to your healthcare provider about registering with the National Pregnancy Registry for Psychiatric Medications by calling 1-866-961-2388 or <a href="http://www.womensmentalhealth.org/peg">www.womensmentalhealth.org/peg</a>.</li> </ul> </li> <li>• are breastfeeding or plan to breastfeed. It is not known if Qelbree passes into breast milk. Talk to your healthcare provider before breastfeeding.</li> <li>• <b>Tell your healthcare provider about all the medicines you or your child takes</b>, including prescription and over-the-counter medicines, vitamins, and herbal supplements. See <b>Do not take Qelbree if you or your child:</b>.</li> </ul> <p>Know the medicines you or your child takes. Keep a list of them and show it to your healthcare provider and pharmacist when you or your child gets a new medicine.</p> <p><b>Do not start any new medicines during treatment with Qelbree without first talking to your healthcare provider.</b></p>
<p><b>What are the most common side effects of QELBREE?</b></p>	<p><b>The most common side effects of Qelbree in children 6-17 years of age include:</b>  •Sleepiness •tiredness •vomiting •irritability •decreased appetite •nausea •trouble sleeping</p> <p><b>The most common side effects of Qelbree in adults include:</b> •insomnia •headache •sleepiness •tiredness •nausea •decreased appetite •dry mouth •constipation</p> <p><b>Effects on weight.</b> Your healthcare provider should check your or your child’s weight before starting and during treatment with Qelbree.</p> <p>These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.</p>
<p><b>How should I take QELBREE?</b></p>	<ul style="list-style-type: none"> <li>• Take Qelbree exactly as your healthcare provider tells you to take it.</li> <li>• Take Qelbree 1 time each day with or without food •Swallow Qelbree capsules whole. •Do not cut, crush, or chew the capsules.</li> <li>• If Qelbree capsules cannot be swallowed whole, the capsule may be opened, and •the entire contents sprinkled onto a spoonful of pudding or applesauce •take all the food mixture <b>without chewing</b> within 15 minutes for pudding or within 2 hours for applesauce •do not store for later use</li> <li>• If you or your child misses a dose, talk to your healthcare provider about what you should do. •Do not take Qelbree for a condition for which it was not prescribed. •Do not give Qelbree to other people, even if they have the same symptoms you have.</li> </ul>
<p>If you or your child takes too much Qelbree or overdoses, call your poison control center at 1-800-222-1222 right away or go to the nearest emergency room.</p> <p><b>Keep Qelbree and all medicines out of the reach of children.</b></p>	
<p>This is not all the information for Qelbree. See the full prescribing information, including boxed warning, at <a href="http://www.Qelbree.com">www.Qelbree.com</a></p>	