

For the treatment of attention-deficit/
hyperactivity disorder (ADHD) in adults...


What are some
reasonable
expectations for treatment
with a nonstimulant?

When can I expect results?

Qelbree is a first-of-its-kind, once-a-day nonstimulant treatment approach for ADHD

- **Qelbree** is the first novel nonstimulant approved for adult ADHD in over 20 years!
- ADHD is with you every day and **Qelbree** is taken once a day, every day

Qelbree is proven to work in medical studies that were conducted over 6-8 weeks—showing significant reduction in ADHD symptom scores in 4 medical studies of more than 1400 people



- **Qelbree** reduced ADHD symptom scores as early as week 2 for some adults

Not everyone responds to medication the same way; your body may need time to adapt. It is important to share how you are feeling with your healthcare provider (HCP), so treatment can be adjusted if necessary.

IMPORTANT SAFETY INFORMATION

Qelbree may increase suicidal thoughts and actions, in children and adults with ADHD, especially within the first few months of treatment or when the dose is changed. Tell your doctor if you or your child have (or if there is a family history of) suicidal thoughts or actions before starting Qelbree. Monitor your or your child's moods, behaviors, thoughts, and feelings during treatment with Qelbree. Report any new or sudden changes in these symptoms right away.

Please see full Important Safety Information on page 4.

What should I know about safety?

Qelbree has been proven safe* in the adult medical study with manageable side effects when taken as directed

- 9% of participants taking **Qelbree** discontinued the medical study due to side effects vs 5% taking inactive treatment (placebo)
- **Qelbree** is unlikely to have negative interactions with stimulants like methylphenidate or amphetamine

*Important: Monitor for mood or behavior changes. Your HCP should also monitor for changes in blood pressure and heart rate

Qelbree is a nonstimulant, which means it is not a narcotic or a controlled medication

- There is no evidence of abuse

Take Qelbree every day. It may take some time for your HCP to optimize your dose. Be sure to share how you are feeling, including any side effects, so your treatment can be adjusted if necessary.

IMPORTANT SAFETY INFORMATION

The most common side effects of **Qelbree** in patients 6 to 17 years are sleepiness, not feeling hungry, feeling tired, nausea, vomiting, trouble sleeping, and irritability, and in adults, insomnia, headache, sleepiness, tiredness, nausea, decreased appetite, dry mouth, and constipation. These are not all the possible side effects of Qelbree.



Can I expect to find it at the pharmacy?

Qelbree is widely available in pharmacies

Patients that were prescribed Qelbree came from

65% switched to Qelbree from a controlled substance or stimulant medication

35% switched to Qelbree from another nonstimulant medication

Qelbree is backed by a robust savings and support program



- Commercially insured patients may pay as little as \$20 per prescription*



Scan here
for savings!

*Terms and conditions apply.

If you are having difficulty getting your ADHD prescription, ask your HCP if Qelbree is an appropriate treatment option. Your HCP may have samples of Qelbree or can request them from the manufacturer if it's decided a trial of Qelbree is an option for you.

IMPORTANT SAFETY INFORMATION

You should not take Qelbree if you or your child:

take a medicine for depression called a monoamine oxidase inhibitor (MAOI) or have stopped taking an MAOI in the past 14 days. Also, you or your child should avoid alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, and theophylline.

Please see full Important Safety Information on page 4.

What else can I expect from treatment?



Qelbree is convenient

- Qelbree is a once-a-day treatment that offers full-day medication coverage
- Qelbree can be prescribed by your doctor via telehealth; there is no need for a new prescription each month to refill your medication



Qelbree may offer you consistency

- Qelbree is taken once a day—any time of day
- Qelbree offers rapid and extended-release, full-day medication coverage
- Qelbree is with you (and your ADHD symptoms) throughout your day



To hear from real patients about their experience on Qelbree visit Qelbree.com.

IMPORTANT SAFETY INFORMATION

Qelbree can increase blood pressure and heart rate. Your or your child's doctor will monitor these vital signs.

Qelbree[®]
ONCE-A-DAY
viloxazine
extended-release capsules
100 mg 150 mg 200 mg

INDICATION

Qelbree is a prescription medicine used to treat ADHD in adults and children 6 years and older.

IMPORTANT SAFETY INFORMATION

Qelbree may increase suicidal thoughts and actions, in children and adults with ADHD, especially within the first few months of treatment or when the dose is changed. Tell your doctor if you or your child have (or if there is a family history of) suicidal thoughts or actions before starting Qelbree. Monitor your or your child's moods, behaviors, thoughts, and feelings during treatment with Qelbree. Report any new or sudden changes in these symptoms right away.

You should not take Qelbree if you or your child:

Take a medicine for depression called a monoamine oxidase inhibitor (MAOI) or have stopped taking an MAOI in the past 14 days. Also, you or your child should avoid alosetron, duloxetine, ramelteon, tasimelteon, tizanidine, and theophylline.

Qelbree can increase blood pressure and heart rate. Your or your child's doctor will monitor these vital signs.

Qelbree may cause manic episodes in patients with bipolar disorder. Tell your doctor if you or your child show any signs of mania.

Do not drive or operate heavy machinery until you know how Qelbree will affect you or your child. Qelbree may cause you or your child to feel sleepy or tired.

The most common side effects of Qelbree in patients 6 to 17 years are sleepiness, not feeling hungry, feeling tired, nausea, vomiting, trouble sleeping, and irritability, and in adults, insomnia, headache, sleepiness, tiredness, nausea, decreased appetite, dry mouth, and constipation. These are not all the possible side effects of Qelbree.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.


Please see Brief Summary, including Boxed Warning, on pages 6 and 7.



Connect with the **Qelbree** Community



Scan here for a conversation starter that can help you focus on key topics to discuss with your HCP.



INDICATION

Qelbree is a prescription medicine used to treat ADHD in patients 6 years and older.

IMPORTANT SAFETY INFORMATION

Qelbree may increase suicidal thoughts and actions, in children and adults with ADHD, especially within the first few months of treatment or when the dose is changed. Tell your doctor if you or your child have (or if there is a family history of) suicidal thoughts or actions before starting Qelbree. Monitor your or your child's moods, behaviors, thoughts, and feelings during treatment with Qelbree. Report any new or sudden changes in these symptoms right away.

Please see full Important Safety Information on page 4.

<p style="text-align: center;">Consumer Brief Summary</p> <p style="text-align: center;">Qelbree® (Kel'bree) (viloxazine extended-release capsules), for oral use</p>	
What is the most important information I should know about QELBREE?	<p>Qelbree can cause serious side effects, including:</p> <ul style="list-style-type: none"> • Increased risk of suicidal thoughts or actions. Qelbree may increase suicidal thoughts or actions in children and adults with attention deficit hyperactivity disorder (ADHD), especially within the first few months of treatment or when the dose is changed. <p>How can I watch for and try to prevent suicidal thoughts and actions?</p> <ul style="list-style-type: none"> • Pay close attention to any changes, especially new or sudden changes in mood, behavior, thoughts, or feelings or if you or your child develops suicidal thoughts or actions. This is very important when Qelbree treatment is started or when the dose is changed. Contact your healthcare provider right away if you or your child has any sudden changes in mood, behavior, thoughts, or feelings, or develops suicidal thoughts or actions. • Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms. <p>Call your healthcare provider or get emergency help right away if you or your child has any of the following symptoms, especially if they are new, worse, or worry you:</p> <ul style="list-style-type: none"> • attempts to commit suicide • new or worse depression • feeling very agitated or restless • trouble sleeping (insomnia) • acting aggressive, being angry or violent • an extreme increase in activity and talking (mania) • thoughts about suicide or dying • new or worse anxiety • panic attacks • new or worse irritability • acting on dangerous impulses • other unusual changes in behavior or mood. <p>See What are the possible side effects of Qelbree? for more information.</p>
What is QELBREE used for?	<p>Qelbree is a prescription medicine used to treat ADHD in adults and children 6 years and older.</p> <p>It is not known if Qelbree is safe and effective in children less than 6 years of age.</p>
Do not take QELBREE if you or your child:	<ul style="list-style-type: none"> • Takes a medicine used to treat depression called a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you or your child takes an MAOI. • Stopped taking an MAOI in the last 14 days. • Takes alossetron, duloxetine, ramelteon, tasimelteon, tizanidine, or theophylline.
What are the possible side effects of QELBREE?	<p>Qelbree can cause serious side effects, including:</p> <ul style="list-style-type: none"> • See What is the most important information I should know about Qelbree? • Increased blood pressure and heart rate. Your healthcare provider should check you or your child's blood pressure and heart rate before starting and during treatment with Qelbree. • Manic episodes. Manic episodes may happen in people with bipolar disorder who take Qelbree. Symptoms may include: •greatly increased energy •racing thoughts •unusually grand ideas •talking more or faster than usual •severe trouble sleeping •reckless behavior •excessive happiness or irritability • Sleepiness and tiredness: Qelbree may cause you or your child to feel sleepy or tired. Do not drive or operate heavy machinery until you know how Qelbree will affect you.
What should I tell my healthcare provider?	Before taking Qelbree, tell your healthcare provider about all your or your child's medical conditions, including if you or your child:

- have a family history of suicide, bipolar disorder, depression, mania or hypomania
- have blood pressure or heart rate problems
- have severe kidney problems. Your healthcare provider may lower the dose of Qelbree.
- are pregnant or plan to become pregnant. Qelbree may cause harm to the mother when taken during pregnancy.
 - Tell your healthcare provider right away if you or your child becomes pregnant or thinks they are pregnant during treatment with Qelbree.
 - There is a pregnancy registry for females who are exposed to Qelbree during pregnancy. Talk to your healthcare provider about registering with the National Pregnancy Registry for Psychiatric Medications by calling 1-866-961-2388 or www.womensmentalhealth.org/preg.
- are breastfeeding or plan to breastfeed. It is not known if Qelbree passes into breast milk. Talk to your healthcare provider before breastfeeding.
- Tell your healthcare provider about all the medicines you or your child takes**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. See **Do not take Qelbree if you or your child**:

Know the medicines you or your child takes. Keep a list of them and show it to your healthcare provider and pharmacist when you or your child gets a new medicine.

Do not start any new medicines during treatment with Qelbree without first talking to your healthcare provider.

What are the most common side effects of QELBREE?

The most common side effects of Qelbree in children 6-17 years of age include:

- Sleepiness •tiredness •vomiting •irritability •decreased appetite •nausea •trouble sleeping

The most common side effects of Qelbree in adults include: •insomnia •headache •sleepiness •tiredness •nausea •decreased appetite •dry mouth •constipation

Effects on weight. Your healthcare provider should check your or your child's weight before starting and during treatment with Qelbree.

These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I take QELBREE?

- Take Qelbree exactly as your healthcare provider tells you to take it.
- Take Qelbree 1 time each day with or without food •Swallow Qelbree capsules whole. •Do not cut, crush, or chew the capsules.
- If Qelbree capsules cannot be swallowed whole, the capsule may be opened, and •the entire contents sprinkled onto a spoonful of pudding or applesauce •take all the food mixture **without chewing** within 15 minutes for pudding or within 2 hours for applesauce •do not store for later use
- If you or your child misses a dose, talk to your healthcare provider about what you should do. •Do not take Qelbree for a condition for which it was not prescribed. •Do not give Qelbree to other people, even if they have the same symptoms you have.

If you or your child takes too much Qelbree or overdoses, call your poison control center at 1-800-222-1222 right away or go to the nearest emergency room.

Keep Qelbree and all medicines out of the reach of children.

This is not all the information for Qelbree. See the full prescribing information, including boxed warning, at www.Qelbree.com